

The head-up cardiopulmonary resuscitation method: Improving neurological outcomes

Anastasiia Bondarenko^{ORCID}, Alla Navolokina^{ORCID}, Marko Kozyk^{ORCID}

European School of Medicine, International European University, Kyiv, Ukraine

The coronavirus disease 2019 (COVID-19) pandemic has significantly impacted the burden on the health service and its efficiency [1]. The pandemic contributed both to the increase in the rate of cardiovascular events as complications of the COVID-19 disease and to the issue of resuscitation and securing the respiratory tract due to the risk of infections [2–4]. For decades, the main form of management of patients experiencing cardiac arrest has been flat or supine cardiopulmonary resuscitation (CPR). No other technique has had a more significant impact on survival than conventional CPR, with the exception of fast defibrillation in appropriate situations. The novel head-up position method of CPR that incorporates controlled elevation of the head and thorax along with active compression decompression and an impedance threshold device has been shown to reduce intracranial pressure and enhance cerebral blood flow, cerebral perfusion pressure, and neurologically favorable survival. When tested on humans, the prefrontal region's median increase in cerebral blood flow in the head-up position was 14.6% higher than in the supine position and this was observed in 83.3% of the patients who were part of the analysis [5]. The main mechanism of benefit for head-up CPR is the use of gravity to improve venous drainage from the paravertebral venous plexus as well as the brain and cerebral venous sinuses, lowering intracranial pressure and opening the possibility for forward blood flow. Heads-up CPR also offers rebalancing blood flow through the lungs similar in manner to patients with heart failure sitting up straight. Another benefit is that compression during head-up CPR compared to conventional CPR is that it lowers

the pressure that is conveyed to the brain via the venous and arterial vasculature, thus preventing concussive damage [6, 7]. Numerous research studies have already proved this in-depth on animal models, but now some demonstrate its efficacy in people [8]. In human models when compared to conventional CPR, the survival rates for the vast majority of out-of-hospital cardiac arrest patients with non-shockable presentations were significantly improved with the head-up position method CPR. Additionally, shorter reaction times to the start of head-up position CPR increased survival odds within highly attainable response times [9]. A significant advantage of this method is the use of automatic chest compression devices, which also relieves the staff by allowing a limited number of people conducting CPR, especially in intensive care units with a large number of COVID-19 patients requiring emergency interventions [10]. Currently, there remains a lack of data on the new resuscitation technique and more extensive research on its use is needed, but the initial results seem convincing and will continue. Its use may be limited by the costs associated with the purchase of new devices — both for automatic chest compression, which are still lacking in many facilities around the world, and for head-up/torso-up positioning devices themselves, however, they can significantly minimize fatigue for emergency medical services personnel in the case of out-of-hospital cardiac arrest and increase the capacity of the hospital systems while significantly improving neurological outcomes. The aforementioned changes could benefit the entire healthcare system.

Conflict of interest: None declared

Address for correspondence: Dr. Alla Navolokina, European School of Medicine, International European University, Akademika Hlushkova Ave, 42B, Kyiv, Ukraine, tel: +380673772377, e-mail: allanavolokina@ieu.edu.pl

Received: 22.11.2022

Accepted: 12.05.2023

Early publication date: 26.05.2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

References

1. Smereka J, Szarpak L, Filipiak K. Modern medicine in COVID-19 era. *Disaster Emerg Med J.* 2020, doi: [10.5603/demj.a2020.0012](https://doi.org/10.5603/demj.a2020.0012).
2. Szarpak L, Filipiak KJ, Skwarek A, et al. Outcomes and mortality associated with atrial arrhythmias among patients hospitalized with COVID-19: A systematic review and meta-analysis. *Cardiol J.* 2022; 29(1): 33–43, doi: [10.5603/CJ.a2021.0167](https://doi.org/10.5603/CJ.a2021.0167), indexed in Pubmed: [34897631](https://pubmed.ncbi.nlm.nih.gov/34897631/).
3. Attila K, Ludwin K, Evrin T, et al. The impact of COVID-19 on airway management in prehospital resuscitation. *Disaster Emerg Med J.* 2020; 5(4): 216–217, doi: [10.5603/demj.a2020.0047](https://doi.org/10.5603/demj.a2020.0047).
4. Nucera G, Chirico F, Rafique Z, et al. Need to update cardiologic guidelines to prevent COVID-19 related myocardial infarction and ischemic stroke. *Cardiol J.* 2022; 29(1): 174–175, doi: [10.5603/CJ.a2021.0120](https://doi.org/10.5603/CJ.a2021.0120), indexed in Pubmed: [34642925](https://pubmed.ncbi.nlm.nih.gov/34642925/).
5. Kim DW, Choi JK, Won SH, et al. A new variant position of head-up CPR may be associated with improvement in the measurements of cranial near-infrared spectroscopy suggestive of an increase in cerebral blood flow in non-traumatic out-of-hospital cardiac arrest patients: A prospective interventional pilot study. *Resuscitation.* 2022; 175: 159–166, doi: [10.1016/j.resuscitation.2022.03.032](https://doi.org/10.1016/j.resuscitation.2022.03.032), indexed in Pubmed: [35395338](https://pubmed.ncbi.nlm.nih.gov/35395338/).
6. Debaty G, Shin SDo, Metzger A, et al. Tilting for perfusion: head-up position during cardiopulmonary resuscitation improves brain flow in a porcine model of cardiac arrest. *Resuscitation.* 2015; 87: 38–43, doi: [10.1016/j.resuscitation.2014.11.019](https://doi.org/10.1016/j.resuscitation.2014.11.019), indexed in Pubmed: [25447353](https://pubmed.ncbi.nlm.nih.gov/25447353/).
7. Moore JC, Segal N, Lick MC, et al. Head and thorax elevation during active compression decompression cardiopulmonary resuscitation with an impedance threshold device improves cerebral perfusion in a swine model of prolonged cardiac arrest. *Resuscitation.* 2017; 121: 195–200, doi: [10.1016/j.resuscitation.2017.07.033](https://doi.org/10.1016/j.resuscitation.2017.07.033), indexed in Pubmed: [28827197](https://pubmed.ncbi.nlm.nih.gov/28827197/).
8. Huang CC, Chen KC, Lin ZY, et al. The effect of the head-up position on cardiopulmonary resuscitation: a systematic review and meta-analysis. *Crit Care.* 2021; 25(1): 376, doi: [10.1186/s13054-021-03797-x](https://doi.org/10.1186/s13054-021-03797-x), indexed in Pubmed: [34717715](https://pubmed.ncbi.nlm.nih.gov/34717715/).
9. Pepe P, Moore J, Bachista K, et al. 3 clinical confirmation of improved likelihood of survival associated with the use of the head-up CPR bundle for non-shockable cardiac arrest presentations. *Ann Emergency Med.* 2022; 80(4): S2, doi: [10.1016/j.annemergmed.2022.08.025](https://doi.org/10.1016/j.annemergmed.2022.08.025).
10. Al-Jeabory M, Borkowska G, Olecka A, et al. Mechanical chest compression devices as an option for out-of-hospital cardiac arrest in COVID-19 pandemic. *Disaster Emerg Med J.* 2021; 6(1): 50–51, doi: [10.5603/demj.a2021.0003](https://doi.org/10.5603/demj.a2021.0003).